

Measure ID	Measure Description	Developer	NQF #
DCR 1	Diabetes: Hemoglobin A1c (HbA1c) Poor Control (>9%): Percentage of patients 18-75 years of age with diabetes who had hemoglobin A1c>9.0% during the measurement period	NCQA	0059
DCR 2	Coronary Artery Disease (CAD): Angiotensin-Converting Enzyme (ACE) Inhibitor or Angiotensin Receptor Blocker (ARB) Therapy-Diabetes or Left Ventricular Systolic Dysfunction (LVEF < 40%): Percentage of patients aged 18 years and older with a diagnosis of coronary artery disease and diabetes seen within a 12-month period who have a current or prior LVEF < 40% who were prescribed an ACE inhibitor or ARB therapy	ACCF	
DCR 3	Diabetes: Medical Attention for Nephropathy: Percentage of patients 18-75 years of age with diabetes who had nephropathy screening test or evidence of nephropathy during the measurement period	NCQA	0062
DCR 4	Tobacco Use: Screening and Cessation Intervention: Percentage of patients aged 18 years and older with diabetes who were screened for tobacco use one or more times within 24 months AND who received cessation counseling intervention if identified as a tobacco user	ACCF/AHA/PCPI	
DCR 5	Dietary Intake Counseling: Percentage of patients aged 18 and older with diabetes who were counseled to eat a healthy diet	ACCF/AHA	
DCR 6	Physical Activity Counseling: Percentage of patients aged 18 and older with diabetes who were counseled to engage in regular physical activity	ACCF/AHA	
DCR 7	Peripheral Artery Disease (PAD): Treatment of Blood Cholesterol to Reduce Atherosclerotic Risk: Percentage of patients 18-75 years of age with PAD and diabetes who were offered moderate-to-high intensity statin	ACCF/AHA/PCPI	
DCR 8	Diabetes Eye Exam (retinal) Performed: Percentage of patients aged 18 and older with diabetes who received an eye exam	NCQA	0055
DCR 9	Diabetes Foot Exam: Percentage of patients aged 18 and older with diabetes who received a foot exam	NCQA	0056
DCR 10	Diabetes Advance Care Plan: Percentage of patients aged 65 years and older with diabetes and who have an advance care plan or surrogate decision marker documented in the medical record or documentation in the medical record that an advance care plan was discussed but the patient did not wish or was not able to name a surrogate decision maker or provide an advance care plan	ACCF/AHA/PCPI	
DCR 11	Antiplatelet Therapy: Percentage of patients aged 18 and older with diabetes who received antiplatelet therapy to reduce the risk of myocardial infarction, stroke, or vascular death in patients with a history of symptomatic PAD	ACCF/AHA/ ACR/SCAI/ SIR/ SVM/ SVN/SVS	

Measure ID	Measure Description	Developer	NQF #
DCR 13	Coronary Artery Disease (CAD): Treatment of Blood Cholesterol to Reduce Atherosclerotic Cardiovascular Risk: Percentage of patients ages 18-75 with CAD and diabetes who were offered moderate-to-high intensity statin	ACCF/AHA/ AMA-PCPI	
DCR 15	Hypertension: Blood Pressure Treatment and Control for High-Risk Patients: Proportion of adults ≥ 18 years of age with both hypertension and diabetes who were prescribed antihypertensive medication or who had adequately controlled blood pressure	ACCF	

Diabetes Collaborative Registry participants include clinicians in primary care, endocrinology, and cardiology ambulatory care settings. Eligible providers must be able to report 7 of the measures listed above to participate in the Diabetes Collaborative Registry.



Diabetes Collaborative Registry
is a Veradigm Network solution

Veradigm® is a business unit of Allscripts©2022 Allscripts Healthcare LLC, and/or its affiliates. All rights reserved. Cited marks are the property of Allscripts Healthcare, LLC and or its affiliates. All other product or company names are the property of their respective holders, all rights reserved.